

Initiating Antiretroviral Therapy in Treatment-Naive Patients

(Updated January 10, 2011)

Panel's Recommendations:

- *Antiretroviral therapy (ART) should be initiated in all patients with a history of an AIDS-defining illness or with a CD4 count <350 cells/mm³ (AI).*
- *ART is also recommended for patients with CD4 counts between 350 and 500 cells/mm³ (A/B*-II).*
- *ART should be initiated, regardless of CD4 count, in patients with the following conditions: HIV-associated nephropathy (HIVAN) (AII) and hepatitis B virus (HBV) coinfection when treatment of HBV is indicated (AIII).*
- *A combination antiretroviral (ARV) drug regimen is also recommended for pregnant women who do not meet criteria for treatment with the goal to prevent perinatal transmission (AI).*
- *For patients with CD4 counts >500 cells/mm³, Panel members are evenly divided: 50% favor starting ART at this stage of HIV disease (B); 50% view initiating therapy at this stage as optional (C) (B/C-III).*
- *Patients initiating ART should be willing and able to commit to lifelong treatment and should understand the benefits and risks of therapy and the importance of adherence (AIII). Patients may choose to postpone therapy, and providers, on a case-by-case basis, may elect to defer therapy based on clinical and/or psychosocial factors.*

Rating of Recommendations: A = Strong; B = Moderate; C = Optional

Rating of Evidence: I = data from randomized controlled trials; II = data from well-designed nonrandomized trials or observational cohort studies with long-term clinical outcomes; III = expert opinion

** Panel members are divided on the strength of this recommendation: 55% voted for strong recommendation (A) and 45% voted for moderate recommendation (B).*

The primary goal of ART is to reduce HIV-associated morbidity and mortality. This is best accomplished by using ART to maximally inhibit HIV replication, as measured by consistent plasma HIV RNA (viral load) values below the level of detection using commercially available assays. Additional benefits of ART, supported by accumulating evidence, are reduction in HIV-associated inflammation and its associated complications and reduction in HIV transmission.

Over the past 20 years, the Panel has made several changes to the recommendations on when to start therapy based on prevailing clinical trial and cohort data and therapeutic options available at the time of each revision. The standard procedure for the Panel is to only make recommendations in agreement with two-thirds of the Panel members. This has not been possible for the **When to Start** recommendations in this version of the guidelines. Accordingly, the breakdown of votes is presented for recommendations supported by less than two-thirds of Panel members.

Randomized controlled trials provide evidence supporting the benefit of ART in patients with CD4 counts ≤ 350 cells/mm³. However, such evidence showing benefit for patients with higher CD4 cell counts is not yet available. Based on cumulative observational cohort data demonstrating benefits of ART in reducing AIDS- and non-AIDS-associated morbidity and mortality, the Panel now recommends ART for patients with CD4 count between 350 and 500 cells/mm³ (**A-B/II**). For patients with CD4 count >500 cells/mm³, Panel members are evenly divided: 50% favor starting ART at earlier stages of HIV disease (**BIII**); 50% view initiating therapy at this stage as optional (**CIII**).

Panel members favoring earlier initiation of therapy base their recommendation on several recent developments: (1) report from at least one recent cohort study demonstrating survival benefit with initiation of ART at CD4 count >500 cells/mm³; (2) growing awareness that untreated HIV infection may be associated with development of many non-AIDS-defining diseases, including cardiovascular disease, kidney disease, liver disease, and malignancy; (3) availability of ARV regimens that are more effective, more convenient, and better tolerated than ARV combinations no longer in use; and (4) increasing evidence that effective ART reduces HIV transmission (**BIII**).

The other 50% of the Panel members feel that current evidence does not definitively demonstrate clear benefit of ART in all patients with CD4 count >500 cells/mm³. They also feel that risks of short- or long-term drug-related complications, nonadherence to lifelong therapy in asymptomatic patients, and potential for development of drug

resistance may offset possible benefits of earlier initiation of therapy. Thus, pending more definitive supporting evidence, these Panel members recommend that therapy in this setting should be optional and considered on a case-by-case basis **(CIII)**.

The known benefits, risks, and limitations of ART, as well as the strength of the recommendations according to CD4 count levels, are discussed below.

BENEFITS OF ANTIRETROVIRAL THERAPY

Earlier studies definitively showed that potent combination ART improves survival and reduces AIDS-related complications in patients with advanced HIV disease. There is now increasing evidence demonstrating the benefits of viral suppression and immunologic responses on reducing mortality and non-AIDS-related complications in patients with higher pretreatment CD4 counts. The following is a focused discussion of the rationale that forms the basis for the Panel's recommendation favoring earlier treatment.

Reduction in Mortality and/or AIDS-Related Morbidity

Patients with a history of an AIDS-defining illness or CD4 count <350 cells/mm³

HIV-infected patients with CD4 counts <200 cells/mm³ are at higher risk of opportunistic diseases, non-AIDS morbidity, and death. Randomized controlled trials in patients with CD4 counts <200 cells/mm³ and/or a history of an AIDS-defining condition provide strong evidence that ART improves survival and delays disease progression in these patients [1-3]. Long-term data from multiple observational cohort studies evaluating earlier ART (>200 cells/mm³) compared with later treatment (<200 cells/mm³) have also provided strong support for these findings [4-8].

Few large, randomized controlled trials address when to start therapy in patients with CD4 counts >200 cells/mm³. CIPRA HT-001 is a randomized clinical trial conducted in Haiti. Study participants were randomized to start ART at CD4 counts of 200–350 cells/mm³ or to defer treatment until their CD4 counts dropped below 200 cells/mm³ or they developed an AIDS-defining condition. In an interim analysis of the study, a higher mortality rate (hazard ratio [HR] = 4.0, $p = 0.0011$) and greater incident tuberculosis (HR = 2.0, $p = 0.0125$) were observed among patients who deferred therapy compared with participants who began ART with CD4 counts of 200–350 cells/mm³ [9]. This evidence led to the study Data Safety Monitoring Board's recommendation to terminate the trial before completion.

The SMART study was a multinational trial enrolling more than 5,400 participants with CD4 counts >350 cells/mm³. Participants were randomized to continuous ART or to treatment interruption until CD4 count dropped to <250 cells/mm³. In a subgroup analysis involving the 249 study participants who were ART naïve at enrollment, a trend of lower risk of serious AIDS- and non-AIDS-related events was seen in those who initiated therapy immediately compared with those who deferred therapy until CD4 count dropped to <250 cells/mm³ ($p = 0.06$) [10].

Collectively, these studies support the Panel's recommendation that ART should be initiated in patients with a history of an AIDS-defining illness or with a CD4 count <350 cells/mm³ **(AI)**.

Patients with a CD4 count between 350 and 500 cells/mm³

There are no randomized trials using current combination regimens in patients with CD4 counts >350 cells/mm³ to provide data that directly address the question of when to start therapy in patients with CD4 counts of 350–500 cells/mm³. Data from the ART Cohort Collaboration (ART-CC), which included 61,798 patient-years of follow-up, showed a declining risk of AIDS or death for up to 5 years in subjects starting therapy with a CD4 count ≥350 cells/mm³ compared with subjects starting between 200 and 349 cells/mm³ [11]. A more recent rigorous analysis of this cohort found that deferring therapy until the 251 to 350 cells/mm³ range was associated with a higher rate of progression to AIDS and death compared with initiating therapy in the 351 to 450 cells/mm³ range (risk ratio: 1.28, 95% confidence interval [CI]: 1.04 to 1.57) [6].

In a collaboration of North American cohort studies (NA-ACCORD) that evaluated patients regardless of whether they had started therapy, the 6,278 patients who deferred therapy until CD4 counts were <350 cells/mm³ had an increased risk of death compared with 2,084 patients who initiated therapy with CD4 counts between 351 and 500 cells/mm³ (risk ratio: 1.69, 95% CI: 1.26 to 2.26) after adjustment for other factors that differed between these two groups [12].

When interpreting both of these cohort studies it is important to note that although the relative risk of a mortality event is evident, the overall number of events was small. In these cohort studies, the relative risks determined could have been influenced by unmeasured confounders that cannot be adjusted for in the analysis. The findings from these observational cohort studies point to potential harm if therapy is deferred until CD4 count falls to <350 cells/mm³. Based on these findings, combined with emerging biologic evidence regarding potential damage to end organs from inflammation associated with untreated HIV replication and the potential reduction in HIV transmission with treatment (see below), the Panel recommends initiation of ART in patients with CD4 counts between 350 and 500 cells/mm³. Panel members are divided on the strength of this recommendation: 55% voted for strong recommendation (A) and 45% voted for moderate recommendation (B) (**A/B-II**).

Patients with a CD4 count >500 cells/mm³

The NA-ACCORD study also observed patients who started treatment at CD4 counts >500 cells/mm³ or after CD4 counts dropped below this threshold. The adjusted mortality rates were significantly higher among the 6,935 patients who deferred therapy until CD4 count fell to <500 cells/mm³ compared with rates in the 2,200 patients who started therapy while CD4 count was >500 cells/mm³ (risk ratio: 1.94, 95% CI: 1.37 to 2.79) [12]. Although large and generally representative of care in the United States, the study has several limitations, including the small number of deaths and the potential for unmeasured confounders that might have influenced outcomes independent of ART.

In contrast, analysis of the ART-CC cohort failed to identify a benefit for patients initiating ART with CD4 counts >450 cells/mm³. This analysis also did not identify a harmful effect of this strategy [6]. Deferral of therapy to the 351–450 cells/mm³ range was associated with a similar rate of progression to AIDS/death compared with initiation of therapy in the 451–550 cells/mm³ range (risk ratio: 0.99, 95% CI: 0.76 to 1.29). This study also found that the proportion of patients with CD4 counts between 451 and 550 cells/mm³ who would progress to AIDS or death before having a CD4 count <450 cells/mm³ was low (1.6%; 95% CI: 1.1 to 2.1%).

Based on these data, along with a better understanding of the pathogenesis of HIV infection and the growing awareness that untreated HIV infection increases the risk of many non-AIDS-defining diseases (see below), 50% of Panel members favor initiation of ART in HIV-infected persons with a CD4 count >500 cells/mm³ (**BIII**).

The other 50% of the Panel members are reluctant to broadly recommend starting ART at higher CD4 cell counts and consider that therapy should be optional at this stage of HIV disease (**CIII**). In making this recommendation, the Panel members note that the amount of data supporting initiation of therapy decreases as the CD4 count increases above 350–500 cells/mm³, and that concerns remain over the unknown overall benefit and long-term risks with earlier treatment.

When discussing starting ART at higher CD4 cell counts (>500 cells/mm³), clinicians should inform patients that data on the clinical benefit of starting treatment at such levels is not conclusive. There is a need for further ongoing research (both with randomized clinical trials and cohort studies) to assess the short- and long-term clinical and public health benefits and cost effectiveness of starting therapy at higher CD4 counts. Such research findings will provide guidance for future recommendations by the Panel.

Effects of Antiretroviral Therapy on HIV-Related Morbidity

HIV-related morbidity and mortality derive not only from immune deficiency but also from direct effects of HIV on specific end organs and the indirect effects of HIV-associated inflammation on these organs. In general, the available data demonstrate that:

- Untreated HIV infection may have detrimental effects at all stages of infection.
- Treatment is beneficial even when initiated later in infection. However, later therapy may not repair damage associated with viral replication during early stages of infection.

- Earlier treatment may prevent the damage associated with HIV replication during early stages of infection.

Clinical studies have demonstrated that sustaining viral suppression and maintaining higher CD4 count, mostly as a result of effective combination ART, delay or prevent some non-AIDS-defining complications, such as HIV-associated kidney disease. Sustained viral suppression and immune recovery may also delay or prevent other disorders, such as liver disease, cardiovascular disease, and malignancies, as discussed below.

HIV-associated nephropathy (HIVAN)

HIVAN is the most common cause of chronic kidney disease in HIV-infected individuals that may lead to end-stage kidney disease [13]. HIVAN is seen almost exclusively in black patients and can occur at any CD4 count. Ongoing viral replication appears to be directly involved in renal injury [14]. HIVAN is extremely uncommon in virologically suppressed patients [15]. ART in patients with HIVAN has been associated with both preserved renal function and prolonged survival [16-18] and therefore should be started in these patients **(AII)**.

Coinfection with hepatitis B virus (HBV) and/or hepatitis C virus (HCV)

HIV infection is associated with more rapid progression of viral hepatitis-related liver disease, including cirrhosis, end-stage liver disease, hepatocellular carcinoma, and fatal hepatic failure [19-20]. Although the mechanisms of accelerated liver disease in HIV-infected patients have not been fully elucidated, HIV-related immunodeficiency and a direct interaction of HIV with hepatic stellate and Kupffer cells have been implicated [21-24]. ART may attenuate liver disease progression in persons coinfecting with HBV and/or HCV by preserving or restoring immune function and reducing HIV-related immune activation and inflammation [25-27]. ARV drugs active against both HIV and HBV (e.g., tenofovir disoproxil fumarate [TDF], lamivudine [3TC], emtricitabine [FTC]) may also prevent the development of significant liver disease by directly suppressing HBV replication [28-29]. Although ARV drugs do not directly inhibit HCV replication, HCV treatment outcomes may be improved if HIV replication is controlled or if CD4 counts are increased [30]. The presence of chronic viral hepatitis increases the risk of ARV-induced liver injury; however, the majority of coinfecting persons do not develop clinically significant liver injury, particularly those receiving recommended ARV regimens [31-33]. Some studies suggest that the rate of hepatotoxicity is greater in persons with more advanced HIV disease. Nevirapine (NVP) toxicity is a notable exception: the hypersensitivity reaction and associated hepatotoxicity to this drug are more frequent in patients with higher CD4 cell counts [34]. Collectively, these data suggest earlier treatment of HIV infection in persons coinfecting with HBV, and possibly HCV **(CIII)**, may reduce the risk of liver disease progression. Furthermore, ART including drugs active against both HIV and HBV should be started in all patients coinfecting with HBV who are also going to receive HBV treatment **(AIII)**.

Cardiovascular disease

Cardiovascular disease is a major cause of mortality among HIV-infected patients, accounting for a third of serious non-AIDS conditions and at least 10% of deaths among HIV-infected patients [35-36]. Studies link exposure to specific ARV drugs to a higher risk of cardiovascular disease [37-38]. Certain HIV treatment regimens are associated with a more atherogenic lipid profile as assessed by lipoprotein particle size analysis among HIV-infected men compared with uninfected controls [39]. Untreated HIV infection may also be associated with an increased risk of cardiovascular disease. In some cross-sectional studies, patients with HIV have higher levels of markers of inflammation and endothelial dysfunction than HIV-uninfected controls [40-42]. In two randomized trials, markers of inflammation and coagulation increased following treatment interruption [43-44]. One study suggests that ART may improve endothelial function [45].

In the SMART study, the risk of cardiovascular events was greater in participants randomized to CD4-guided treatment interruption compared with participants who received continuous ART [46]. In other studies, ART resulted in marked improvement in parameters associated with cardiovascular diseases, including markers of inflammation (e.g., interleukin 6 [IL-6] and high sensitivity C-reactive protein [hsCRP]) and endothelial dysfunction [41, 45]. There is also a modest association between lower CD4 count while on therapy and short-term risk of cardiovascular disease [42, 47-48]. However, in at least one of these cohorts (the CASCADE study), the link between CD4 count and fatal cardiovascular events was no longer statistically significant when adjusting for plasma HIV RNA level. Collectively, the data linking viremia and endothelial dysfunction and inflammation, the increased risk of cardiovascular events

with treatment interruption, and the association between cardiovascular disease and CD4 cell depletion suggest that early control of HIV replication with ART can be used as a strategy to reduce cardiovascular disease risk **(BIII)**.

Malignancies

Several population-based analyses suggest increased incidence of non-AIDS-associated malignancies during chronic HIV infection. The incidence of non-AIDS malignancy in HIV-infected subjects is higher than in matched HIV-uninfected controls [49]. Large cohort studies of mostly patients receiving ART have reported a consistent link between low CD4 counts ($<350\text{--}500\text{ cells/mm}^3$) and the risk of AIDS- and/or non-AIDS-defining malignancy [42, 47, 50-53]. The ANRS C04 Study demonstrated a statistically significant relative risk of all cancers evaluated (except for anal carcinoma) in patients with CD4 counts $<500\text{ cells/mm}^3$ compared with patients with current CD4 counts $>500\text{ cells/mm}^3$ and a protective effect of ART for HIV-associated malignancies [50]. This potential effect of HIV-associated immunodeficiency is particularly striking with regard to cancers associated with chronic viral infections (e.g., HBV, HCV, human papilloma virus [HPV], Epstein-Barr virus [EBV], human herpes virus-8 [HHV-8]) [54-55]. Cumulative HIV viremia itself may also be associated with the risk of non-Hodgkin lymphoma and other AIDS-defining malignancies, independent of other factors [53, 56]. Together this evidence suggests that initiating ART to suppress HIV replication and maintain CD4 counts at above $350\text{--}500\text{ cells/mm}^3$ may reduce the risk of both AIDS-defining and non-AIDS-defining malignancies **(CIII)**.

Neurocognitive decline

Early in the HIV epidemic, HIV was identified in brain tissue [57] and assumed to be the cause of AIDS dementia complex [58]. The improvement of AIDS dementia complex symptoms with the use of ART supported this assumption [59-60]. The CASCADE observational cohort reported a dramatic decline in the incidence of HIV-associated dementia from 6.49 per 1,000 person-years (before 1997) to 0.66 per 1,000 person-years (2003–2006), after the widespread use of potent ART [61]. In this cohort, having a current CD4 count $>350\text{ cells/mm}^3$ was associated with the lowest risk of developing HIV-associated dementia. HIV infection has also been associated with a number of less severe neurologic complications, including changes in neuropsychological ability, speed of processing, and everyday functioning [62]. Such syndromes also were predicted by a lower pretherapy CD4 nadir and/or by CD4 count while on therapy [63-64]. Additional clinical data are needed to determine the relative roles of ongoing HIV replication and potential neurotoxicity of ARV agents in the development of neurocognitive dysfunction. Whether early initiation of therapy will prevent HIV-associated neurocognitive dysfunction remains unclear. However, the neurological complications that may accompany uncontrolled HIV replication and CD4 depletion suggest a potential benefit of earlier initiation of ART **(CIII)**.

Age and treatment-related immune reconstitution

The CD4 cell response to ART is an important predictor of short-term and long-term morbidity and mortality. Treatment initiation at an older age is consistently associated with a less robust CD4 count response; starting therapy at a younger age may result in better immunologic and perhaps clinical outcomes [65-67]**(CIII)**.

T-cell activation and inflammation

Early untreated HIV infection is associated with sustained high-level inflammation and T-cell activation [68-70]. The degree of T-cell activation during untreated HIV disease is associated with risk of subsequent disease progression, independent of other factors such as plasma HIV RNA levels and the peripheral CD4 T-cell count [71-72]. ART results in a rapid, but often incomplete, decrease in most markers of HIV-associated immune activation [73-77]. Persistent T-cell activation and/or T-cell dysfunction is particularly evident among patients who delay therapy until later stage disease (CD4 count $<350\text{ cells/mm}^3$) [74, 77-78]. The degree of persistent inflammation during treatment, as represented by the levels of IL-6, may be independently associated with risk of death [44]. Collectively, these observations support earlier use of ART for at least two reasons. First, treatment decreases the level of inflammation and T-cell activation, which may be associated with reduced short-term risk of AIDS- and non-AIDS-related morbidity and mortality [44, 79-80]. Second, because the degree of residual inflammation and/or T-cell dysfunction during ART appears to be higher in patients with lower CD4 cell nadirs [74, 77-78], earlier treatment may result in less

residual immunological perturbations on therapy, and hence less risk for AIDS- and non-AIDS-related complications (CIII).

Prevention of HIV Transmission

Prevention of mother-to-child transmission

Effective ART reduces transmission of HIV. The most dramatic and well-established example of this effect is the use of ART in pregnant women to prevent mother-to-child transmission (MTCT) of HIV. Effective suppression of HIV replication, as reflected in plasma HIV RNA, is a key determinant in reducing perinatal transmission. In the United States, the use of combination ART during pregnancy has reduced the HIV transmission rate from approximately 20%–30% to <2% [81]. Thus, use of combination ARV drug regimens is recommended for all HIV-infected pregnant women to prevent MTCT of HIV (AI), even if the mother does not meet the criteria for initiation of therapy for treatment of her HIV infection. Following delivery, considerations regarding continuation of the ARV regimen for maternal therapeutic indications are the same as for other nonpregnant individuals. For detailed recommendations, see [Perinatal Guidelines](#) [82].

Prevention of sexual transmission

Emerging evidence supports the concept of "treatment as prevention" of sexual transmission of HIV. Lower plasma HIV RNA levels are associated with decreases in the concentration of the virus in genital secretions [83–84]. Studies of HIV serodiscordant heterosexual couples have demonstrated a relationship between the level of plasma viremia and HIV transmission risk: when plasma HIV RNA levels are lower, transmission events are less common [85–89]. These investigations, as well as other observational studies and modeling analyses demonstrating a decreased rate of HIV transmission among serodiscordant heterosexual couples following the introduction of ART, suggest that suppression of viremia in ART-adherent patients with no concomitant sexually transmitted infections (STIs) substantially reduces the risk of HIV transmission [88–93]. Based on these studies, the use of effective ART regardless of CD4 count is likely to reduce transmission to the uninfected sexual partner (BII).

POTENTIAL LIMITATIONS OF EARLIER INITIATION OF THERAPY

Although there are benefits associated with earlier initiation of ART, there are also potential limitations to this approach. Concerns about long-term toxicity and the development of ARV resistance have served as a rationale for the deferral of HIV therapy. Earlier initiation of ART at higher CD4 counts (e.g., >500 cells/mm³) results in greater cumulative time on therapy. Assuming treatment for many decades after initiation, the additional therapy represents a small percentage of the total time on ART for most patients.

Although newer ARV regimens are generally better tolerated, more convenient, and more potent than older regimens, there are fewer longer term safety data for the newer agents. Analyses supporting ART initiation at CD4 counts >350 cells/mm³ (e.g., NA-ACCORD and ART-CC) were conducted with cohorts largely treated with regimens less commonly used in clinical practice. These studies reported on clinical endpoints of death and/or AIDS disease progression but lacked information on drug toxicities, resistance, or adherence. Therefore, in considering earlier initiation of therapy, concerns for some adverse consequences of ART remain.

Antiretroviral Drug Toxicities and Quality of Life

Earlier initiation of ART extends exposure to ARV agents by several years. The D:A:D study found an increased incidence of cardiovascular disease associated with cumulative exposure to some drugs within the nucleoside reverse transcriptase inhibitor (NRTI) and protease inhibitor (PI) classes [38, 94]. In the SMART study, continuous exposure to ART has been associated with significantly greater loss of bone density compared with interruption or deferral of therapy [46]. There may be unknown complications related to cumulative use of ARV drugs for many decades. A list of known ARV-associated toxicities can be found in [Adverse Effects of Antiretroviral Agents](#).

Although ART frequently improves quality of life among symptomatic patients, it may also be associated with reduced quality of life in some patients, especially those who are asymptomatic at initiation of therapy. Although better tolerated and easier to administer than older drugs, most ARV drugs now used in first-line regimens can cause side effects that may reduce quality of life. Efavirenz (EFV), for example, can cause neurocognitive or psychiatric side effects, and all the PIs have been associated with gastrointestinal side effects. Furthermore, some patients may find that the inconvenience of taking medication every day outweighs the overall benefit and might choose to delay therapy whenever possible.

Drug Resistance

Very early treatment initiation may lead to an earlier onset of drug resistance selection in nonadherent patients. The consequent harm is loss of important drugs or drug classes and risk of transmission of drug-resistant HIV. Some asymptomatic patients may be less motivated to remain adherent to their HIV treatment regimens if treatment is initiated far in advance of an immediate risk of HIV-associated morbidity and mortality. The greater convenience and potency of current ARV regimens facilitate adherence and reduce the risk of ARV resistance. One study suggests that the risk of drug resistance at the time of virologic failure is lower among patients who initiated treatment at higher CD4 counts [95]. Treatment adherence is key to viral suppression and should be stressed prior to initiation of therapy and during follow-up visits.

Nonadherence to Antiretroviral Therapy

At any CD4 count, adherence to therapy is essential to achieve viral suppression and prevent emergence of resistance mutations. Several behavioral and social factors associated with poor adherence, such as untreated major psychiatric disorders, active substance abuse, social circumstances, patient concerns about side effects, and poor adherence to clinic visits, have been identified. Clinicians should identify areas where additional intervention is needed to improve adherence both before and after initiation of therapy. Some strategies to improve adherence are discussed in

[Adherence](#).

Cost

Although ART adds to the annual cost of treatment, several modeling studies support the cost effectiveness of HIV therapy initiated soon after diagnosis [96-98]. Studies have reported that the annual cost of care is 2½ times higher for patients with CD4 counts <50 cells/mm³ compared with patients with CD4 counts >350 cells/mm³ [99]. A large proportion of the health care expenditure in patients with advanced infection is from non-ARV drugs and hospitalization. However, no cost comparisons have been reported between those starting ART with a CD4 count between 350 and 500 cells/mm³ versus >500 cells/mm³.

SUMMARY

In earlier versions of these treatment guidelines, concerns about long-term toxicity, reduced quality of life, and the potential for drug resistance served as key reasons to defer HIV therapy for as long as possible. Inherent in this argument was the assumption that the harm associated with viral replication was less than the harm associated with the toxicities of ARV drugs in patients with higher CD4 counts. There is now more evidence that untreated HIV infection has negative consequences on health at all stages of disease. Also, the drug combinations now available are better tolerated than previous regimens, leading to greater efficacy and improved adherence [100]. The current guidelines therefore emphasize avoiding adverse consequences of untreated HIV infection while managing potential drug toxicity.

RECOMMENDATIONS

Based on the cumulative weight of evidence described above, the Panel recommends that:

- ART should be initiated in all patients with a history of an AIDS-defining illness or with a CD4 count of <350 cells/mm³ (**AI**).
- ART is also recommended for patients with CD4 counts between 350 and 500 cells/mm³ (**A/B-II**).
- ART should also be initiated, regardless of CD4 count, in patients with the following conditions: HIVAN (**AI**) and HBV coinfection when treatment of HBV is indicated (**AIII**).
- A combination ARV drug regimen is also recommended for pregnant women who do not meet criteria for treatment with the goal to prevent perinatal transmission (**AI**).
- For patients with CD4 counts >500 cells/mm³, 50% of the Panel members favor starting ART (B); the other 50% of members view treatment as optional (C) in this setting (**B/C-III**).
- Patients initiating ART should be willing and able to commit to lifelong treatment and should understand the benefits and risks of therapy and the importance of adherence (**AIII**).
- Patients may choose to postpone therapy, and providers, on a case-by-case basis, may elect to defer therapy based on clinical and/or psychosocial factors.

* The Panel is divided on the strength of this recommendation: 55% of Panel members voted for strong recommendation (**A**) and 45% voted for moderate recommendation (**B**).

Conditions Favoring More Rapid Initiation of Therapy

Deferring ART may be appropriate in some cases. However, several conditions increase the urgency for therapy, including:

- Pregnancy (**AI**) (Clinicians should refer to the [Perinatal Guidelines](#) for more detailed recommendations for the management of HIV-infected pregnant women.) [82]
- AIDS-defining conditions (**AI**)
- Acute opportunistic infections (see discussion below)
- Lower CD4 counts (e.g., <200 cells/mm³) (**AI**)
- Rapidly declining CD4 counts (e.g., >100 cells/mm³ decrease per year) (**AIII**)
- Higher viral loads (e.g., >100,000 copies/mL) (**BII**)
- HIVAN (**AI**)
- HBV coinfection when treatment for HBV is indicated (**AIII**)

Acute opportunistic infections

In patients with opportunistic conditions for which there is no effective therapy (e.g., cryptosporidiosis, microsporidiosis, progressive multifocal leukoencephalopathy) but for which ART may improve outcomes by improving immune responses, the benefits of ART outweigh any increased risk, and therefore treatment should be started as soon as possible (**AIII**).

In the setting of opportunistic infections, such as cryptococcal meningitis or non-tuberculous mycobacterial infections, for which immediate therapy may increase the risk of immune reconstitution inflammatory syndrome (IRIS), a short delay may be warranted before initiating ART [101-102] (**CIII**).

In the setting of other opportunistic infections, such as *Pneumocystis jiroveci* pneumonia (PCP), early initiation of ART is associated with increased survival, and therapy should not be delayed [3] (**AI**).

In patients who have active tuberculosis, initiating ART within the first 1–2 months of treatment for tuberculosis appears to confer a significant survival advantage [103-104]. (See [Mycobacterium Tuberculosis Disease with HIV Coinfection](#).)

Clinicians should refer to the [Guidelines for Prevention and Treatment of Opportunistic Infections in HIV-Infected Adults and Adolescents](#) [105] for more detailed discussion on when to initiate ART in the setting of a specific opportunistic infection.

Conditions Where Deferral of Therapy Might be Considered

Some patients and their clinicians may decide to defer therapy for a period of time based on clinical or personal circumstances. The degree to which these factors might argue for deferral of therapy depends on the CD4 count and viral load. Although deferring therapy for the reasons discussed below may be reasonable for patients with high CD4 counts (e.g., >500 cells/mm³), deferral for patients with much lower CD4 counts (e.g., <200 cells/mm³) should be considered only in rare situations and should be undertaken with close clinical follow-up. A brief delay in initiating therapy may be considered to allow a patient more time to prepare for lifelong treatment.

When there are significant barriers to adherence

Deferring treatment for patients with higher CD4 counts who are at risk of poor adherence may be prudent while the barriers to adherence are being addressed. However, increased urgency for ART (see above) may override potential predictors of poor adherence.

Several methodologies exist to help providers assess adherence. When the most feasible measure of adherence is self-report, this assessment should be completed at each clinic visit, using one of the available reliable and valid instruments [106-107]. If other objective measures are available (e.g., pharmacy refill data, pill count), these methods should also be implemented as therapy begins [108-110]. Continuous assessment and counseling make it possible for the clinician to intervene early to address barriers to adherence occurring at any point during treatment.

Presence of comorbidities that complicate or prohibit antiretroviral therapy

Deferral of ART may be considered when either the treatment or manifestations of other medical conditions could complicate the treatment of HIV infection or vice versa. Examples include:

- Patients requiring surgery that might result in an extended interruption of ART.
- Patients taking medications that have clinically significant drug interactions with ART and for whom alternative therapy is not available.

In each of these cases, it is assumed that the situation is temporary and that ART will be initiated after the conflicting condition has resolved.

There are some less common situations in which ART may not be indicated at any time while CD4 counts remain high. In particular, such situations include patients with a poor prognosis due to a concomitant medical condition who would not be expected to gain survival or quality-of-life benefits from ART. Examples include patients with incurable non-HIV-related malignancies or end-stage liver disease who are not being considered for liver transplantation. The decision to forego ART in such patients may be easier in those with higher CD4 counts; they are likely asymptomatic for HIV, and their survival is unlikely to be prolonged by ART. However, it should be noted that ART may improve outcomes, including survival, in patients with some HIV-associated malignancies (e.g., lymphoma or Kaposi's sarcoma) and in patients with liver disease due to chronic HBV or HCV.

Elite HIV controllers or long-term nonprogressors

A small subset of ARV-untreated HIV-infected persons (~3%–5%) are able to maintain normal CD4 cell counts for many years (long-term nonprogressors), while an even smaller subset (~1%) are able to maintain suppressed viral loads for years (elite controllers). It is possible that such patients would not benefit from ART. However, some nonprogressors have high viral loads, and some elite controllers progress clinically or immunologically [111-112]. Although therapy may be theoretically beneficial for patients in either group, clinical data supporting therapy for nonprogressors and elite controllers are lacking.

THE NEED FOR EARLY DIAGNOSIS OF HIV

Fundamental to the earlier initiation of therapy recommended in these guidelines is the assumption that patients will be diagnosed early in the course of HIV infection, making earlier initiation of therapy an option. Unfortunately, most HIV-infected patients are not diagnosed until they are at much later stages of disease [113-116]. Despite the 2006 Centers for Disease Control and Prevention (CDC) recommendations for routine, opt-out HIV screening in the health care setting [117] regardless of perceived risk of infection, the median CD4 count for newly diagnosed patients remains in the ~200 cells/mm³ range. (The exception is pregnant women diagnosed during prenatal care, who have a much higher median initial CD4 count.) Delay in HIV diagnosis is more often seen in nonwhites, injection drug users, and older patients; a substantial proportion of these individuals develop AIDS-defining illnesses within 1 year of diagnosis [113-116]. Therefore, for the current treatment guidelines to have maximum impact, routine HIV screening per current CDC recommendations is essential. It is critical that all newly diagnosed patients be educated about HIV disease and linked to care for full evaluation, follow-up, and management. Once in care, focused effort is required to retain patients in the health care system.

CONCLUSION

The current recommendations are based on increasing evidence that supports earlier initiation of ART than was advocated in previous guidelines. The strength of the recommendations varies with the quality and availability of existing evidence. Panel members are divided regarding the strength of recommendations for starting therapy in patients with higher CD4 cell counts, as discussed above. The Panel will continue to monitor and assess the results of ongoing and planned randomized clinical trials and observational studies, which will provide the Panel with additional guidance to form future recommendations.

References

1. HIV Trialists' Collaborative Group. Zidovudine, didanosine, and zalcitabine in the treatment of HIV infection: meta-analyses of the randomised evidence. *Lancet*. 1999;353(9169):2014-2025.
2. Hammer SM, Squires KE, Hughes MD, et al. A controlled trial of two nucleoside analogues plus zidovudine in persons with human immunodeficiency virus infection and CD4 cell counts of 200 per cubic millimeter or less. AIDS Clinical Trials Group 320 Study Team. *N Engl J Med*. 1997;337(11):725-733.
3. Zolopa A, Andersen J, Powderly W, et al. Early antiretroviral therapy reduces AIDS progression/death in individuals with acute opportunistic infections: a multicenter randomized strategy trial. *PLoS One*. 2009;4(5):e5575.
4. Mocroft A, Vella S, Benfield TL, et al. Changing patterns of mortality across Europe in patients infected with HIV-1. EuroSIDA Study Group. *Lancet*. 1998;352(9142):1725-1730.
5. Hogg RS, Yip B, Chan KJ, et al. Rates of disease progression by baseline CD4 cell count and viral load after initiating triple-drug therapy. *JAMA*. 2001;286(20):2568-2577.
6. Sterne JA, May M, Costagliola D, et al. Timing of initiation of antiretroviral therapy in AIDS-free HIV-1-infected patients: a collaborative analysis of 18 HIV cohort studies. *Lancet*. 2009;373(9672):1352-1363.
7. Baker JV, Peng G, Rapkin J, et al. CD4+ count and risk of non-AIDS diseases following initial treatment for HIV infection. *AIDS*. 2008;22(7):841-848.
8. Palella FJ, Jr., Deloria-Knoll M, Chmiel JS, et al. Survival benefit of initiating antiretroviral therapy in HIV-infected persons in different CD4+ cell strata. *Ann Intern Med*. 2003;138(8):620-626.
9. Severe P, Juste MA, Ambroise A, et al. Early versus standard antiretroviral therapy for HIV-infected adults in Haiti. *N Engl J Med*. 2010;363(3):257-265.
10. Emery S, Neuhaus JA, Phillips AN, et al. Major clinical outcomes in antiretroviral therapy (ART)-naïve participants and in those not receiving ART at baseline in the SMART study. *J Infect Dis*. 2008;197(8):1133-1144.
11. May M, Sterne JA, Sabin C, et al. Prognosis of HIV-1-infected patients up to 5 years after initiation of HAART: collaborative analysis of prospective studies. *AIDS*. 2007;21(9):1185-1197.
12. Kitahata MM, Gange SJ, Abraham AG, et al. Effect of early versus deferred antiretroviral therapy for HIV on survival. *N Engl J Med*. 2009;360(18):1815-1826.
13. Szczec LA, Gupta SK, Habash R, et al. The clinical epidemiology and course of the spectrum of renal diseases associated with HIV infection. *Kidney Int*. 2004;66(3):1145-1152.
14. Marras D, Bruggeman LA, Gao F, et al. Replication and compartmentalization of HIV-1 in kidney epithelium of patients with HIV-associated nephropathy. *Nat Med*. 2002;8(5):522-526.
15. Estrella M, Fine DM, Gallant JE, et al. HIV type 1 RNA level as a clinical indicator of renal pathology in HIV-infected patients. *Clin Infect Dis*. 2006;43(3):377-380.

16. Atta MG, Gallant JE, Rahman MH, et al. Antiretroviral therapy in the treatment of HIV-associated nephropathy. *Nephrol Dial Transplant*. 2006;21(10):2809-2813.
17. Schwartz EJ, Szczech LA, Ross MJ, et al. Highly active antiretroviral therapy and the epidemic of HIV+ end-stage renal disease. *J Am Soc Nephrol*. 2005;16(8):2412-2420.
18. Kalayjian RC, Franceschini N, Gupta SK, et al. Suppression of HIV-1 replication by antiretroviral therapy improves renal function in persons with low CD4 cell counts and chronic kidney disease. *AIDS*. 2008;22(4):481-487.
19. Thein HH, Yi Q, Dore GJ, et al. Natural history of hepatitis C virus infection in HIV-infected individuals and the impact of HIV in the era of highly active antiretroviral therapy: a meta-analysis. *AIDS*. 2008;22(15):1979-1991.
20. Thio CL, Seaberg EC, Skolasky R, Jr., et al. HIV-1, hepatitis B virus, and risk of liver-related mortality in the Multicenter Cohort Study (MACS). *Lancet*. 2002;360(9349):1921-1926.
21. Weber R, Sabin CA, Friis-Moller N, et al. Liver-related deaths in persons infected with the human immunodeficiency virus: the D:A:D study. *Arch Intern Med*. 2006;166(15):1632-1641.
22. Balagopal A, Philp FH, Astemborski J, et al. Human immunodeficiency virus-related microbial translocation and progression of hepatitis C. *Gastroenterology*. 2008;135(1):226-233.
23. Blackard JT, Kang M, St Clair JB, et al. Viral factors associated with cytokine expression during HCV/HIV co-infection. *J Interferon Cytokine Res*. 2007;27(4):263-269.
24. Hong F, Tuyama A, Lee TF, et al. Hepatic stellate cells express functional CXCR4: role in stromal cell-derived factor-1alpha-mediated stellate cell activation. *Hepatology*. 2009;49(6):2055-2067.
25. Macias J, Berenguer J, Japon MA, et al. Fast fibrosis progression between repeated liver biopsies in patients coinfecting with human immunodeficiency virus/hepatitis C virus. *Hepatology*. 2009;50(4):1056-1063.
26. Verma S, Goldin RD, Main J. Hepatic steatosis in patients with HIV-Hepatitis C Virus coinfection: is it associated with antiretroviral therapy and more advanced hepatic fibrosis? *BMC Res Notes*. 2008;1:46.
27. Ragni MV, Nalesnik MA, Schillo R, et al. Highly active antiretroviral therapy improves ESLD-free survival in HIV-HCV co-infection. *Haemophilia*. 2009;15(2):552-558.
28. Matthews GV, Avihingsanon A, Lewin SR, et al. A randomized trial of combination hepatitis B therapy in HIV/HBV coinfecting antiretroviral naive individuals in Thailand. *Hepatology*. 2008;48(4):1062-1069.
29. Peters MG, Andersen J, Lynch P, et al. Randomized controlled study of tenofovir and adefovir in chronic hepatitis B virus and HIV infection: ACTG A5127. *Hepatology*. 2006;44(5):1110-1116.
30. Avidan NU, Goldstein D, Rozenberg L, et al. Hepatitis C viral kinetics during treatment with peg IFN-alpha-2b in HIV/HCV coinfecting patients as a function of baseline CD4+ T-cell counts. *J Acquir Immune Defic Syndr*. 2009;52(4):452-458.
31. Clotet B, Bellos N, Molina JM, et al. Efficacy and safety of darunavir-ritonavir at week 48 in treatment-experienced patients with HIV-1 infection in POWER 1 and 2: a pooled subgroup analysis of data from two randomised trials. *Lancet*. 2007;369(9568):1169-1178.
32. Steigbigel RT, Cooper DA, Kumar PN, et al. Raltegravir with optimized background therapy for resistant HIV-1 infection. *N Engl J Med*. 2008;359(4):339-354.
33. Molina JM, Andrade-Villanueva J, Echevarria J, et al. Once-daily atazanavir/ritonavir compared with twice-daily lopinavir/ritonavir, each in combination with tenofovir and emtricitabine, for management of antiretroviral-naïve HIV-1-infected patients: 96-week efficacy and safety results of the CASTLE study. *J Acquir Immune Defic Syndr*. 2010;53(3):323-332.
34. van Leth F, Andrews S, Grinsztejn B, et al. The effect of baseline CD4 cell count and HIV-1 viral load on the efficacy and safety of nevirapine or efavirenz-based first-line HAART. *AIDS*. 2005;19(5):463-471.
35. Smith C. Factors associated with specific causes of death amongst HIV-positive individuals in the D:A:D Study. *AIDS*. 2010;24(10):1537-1548.
36. Mocroft A, Reiss P, Gasiorowski J, et al. Serious fatal and nonfatal non-AIDS-defining illnesses in Europe. *J Acquir Immune Defic Syndr*. 2010;55(2):262-270.
37. Sabin CA, Worm SW, Weber R, et al. Use of nucleoside reverse transcriptase inhibitors and risk of myocardial infarction in HIV-infected patients enrolled in the D:A:D study: a multi-cohort collaboration. *Lancet*. 2008;371(9622):1417-1426.
38. Friis-Moller N, Reiss P, Sabin CA, et al. Class of antiretroviral drugs and the risk of myocardial infarction. *N Engl J Med*. 2007;356(17):1723-1735.
39. Riddler SA, Li X, Otvos J, et al. Antiretroviral therapy is associated with an atherogenic lipoprotein phenotype among HIV-1-infected men in the Multicenter AIDS Cohort Study. *J Acquir Immune Defic Syndr*. 2008;48(3):281-288.
40. Ross AC, Armentrout R, O'Riordan MA, et al. Endothelial activation markers are linked to HIV status and are independent of antiretroviral therapy and lipodystrophy. *J Acquir Immune Defic Syndr*. 2008;49(5):499-506.
41. McComsey G, Smith K, Patel P, et al. Similar reductions in markers of inflammation and endothelial activation after initiation of abacavir/lamivudine or tenofovir/emtricitabine: The HEAT Study. Paper presented at: 16th Conference on Retroviruses and Opportunistic Infections; 2009; Montreal, Canada.
42. Baker JV, Duprez D, Rapkin J, et al. Untreated HIV infection and large and small artery elasticity. *J Acquir Immune Defic Syndr*. 2009;52(1):25-31.
43. Calmy A, Gayet-Ageron A, Montecucco F, et al. HIV increases markers of cardiovascular risk: results from a randomized, treatment interruption trial. *AIDS*. 2009;23(8):929-939.
44. Kuller LH, Tracy R, Belloso W, et al. Inflammatory and coagulation biomarkers and mortality in patients with HIV infection. *PLoS Med*. 2008;5(10):e203.
45. Torriani FJ, Komarow L, Parker RA, et al. Endothelial function in human immunodeficiency virus-infected antiretroviral-naïve subjects before and after starting potent antiretroviral therapy: The ACTG (AIDS Clinical Trials Group) Study 5152s. *J Am Coll Cardiol*. 2008;52(7):569-576.

46. El-Sadr WM, Lundgren JD, Neaton JD, et al. CD4+ count-guided interruption of antiretroviral treatment. *N Engl J Med*. 2006;355(22):2283-2296.
47. Marin B, Thiebaut R, Bucher HC, et al. Non-AIDS-defining deaths and immunodeficiency in the era of combination antiretroviral therapy. *AIDS*. 2009;23(13):1743-1753.
48. Phillips AN, Neaton J, Lundgren JD. The role of HIV in serious diseases other than AIDS. *AIDS*. 2008;22(18):2409-2418.
49. Bedimo RJ, McGinnis KA, Dunlap M, et al. Incidence of non-AIDS-defining malignancies in HIV-infected versus noninfected patients in the HAART era: impact of immunosuppression. *J Acquir Immune Defic Syndr*. 2009.
50. Guiguet M, Boue F, Cadranet J, et al. Effect of immunodeficiency, HIV viral load, and antiretroviral therapy on the risk of individual malignancies (FHDH-ANRS CO4): a prospective cohort study. *Lancet Oncol*. 2009.
51. Monforte A, Abrams D, Pradier C, et al. HIV-induced immunodeficiency and mortality from AIDS-defining and non-AIDS-defining malignancies. *AIDS*. 2008;22(16):2143-2153.
52. Reekie J, Mocroft A, Engsig F, et al. Relationship between current level of immunodeficiency and non-AIDS-defining malignancies. Paper presented at: 16th Conference on Retroviruses and Opportunistic Infections; February, 2009; Montreal, Canada. Abstract 860a.
53. Bruyand M, Thiebaut R, Lawson-Ayayi S, et al. Role of uncontrolled HIV RNA level and immunodeficiency in the occurrence of malignancy in HIV-infected patients during the combination antiretroviral therapy era: Agence Nationale de Recherche sur le Sida (ANRS) CO3 Aquitaine Cohort. *Clin Infect Dis*. 2009;49(7):1109-1116.
54. Silverberg MJ, Chao C, Leyden WA, et al. HIV infection and the risk of cancers with and without a known infectious cause. *AIDS*. 2009;23(17):2337-2345.
55. Grulich AE, van Leeuwen MT, Falster MO, et al. Incidence of cancers in people with HIV/AIDS compared with immunosuppressed transplant recipients: a meta-analysis. *Lancet*. 2007;370(9581):59-67.
56. Zoufaly A, Stellbrink HJ, Heiden MA, et al. Cumulative HIV viremia during highly active antiretroviral therapy is a strong predictor of AIDS-related lymphoma. *J Infect Dis*. 2009;200(1):79-87.
57. Shaw GM, Harper ME, Hahn BH, et al. HTLV-III infection in brains of children and adults with AIDS encephalopathy. *Science*. 1985;227(4683):177-182.
58. Janssen RS, Nwanyanwu OC, Selik RM, et al. Epidemiology of human immunodeficiency virus encephalopathy in the United States. *Neurology*. 1992;42(8):1472-1476.
59. Schmitt FA, Bigley JW, McKinnis R, et al. Neuropsychological outcome of zidovudine (AZT) treatment of patients with AIDS and AIDS-related complex. *N Engl J Med*. 1988;319(24):1573-1578.
60. Robertson KR, Robertson WT, Ford S, et al. Highly active antiretroviral therapy improves neurocognitive functioning. *J Acquir Immune Defic Syndr*. 2004;36(1):562-566.
61. Bhaskaran K, Mussini C, Antinori A, et al. Changes in the incidence and predictors of human immunodeficiency virus-associated dementia in the era of highly active antiretroviral therapy. *Ann Neurol*. 2008;63(2):213-221.
62. Vance D, Wadley V, Crowe M, et al. Cognitive and everyday functioning in younger and older adults with and without HIV (in press). *Clinical Gerontologist*.
63. Robertson KR, Smurzynski M, Parsons TD, et al. The prevalence and incidence of neurocognitive impairment in the HAART era. *AIDS*. 2007;21(14):1915-1921.
64. Munoz-Moreno JA, Fumaz CR, Ferrer MJ, et al. Nadir CD4 cell count predicts neurocognitive impairment in HIV-infected patients. *AIDS Res Hum Retroviruses*. 2008;24(10):1301-1307.
65. The Collaboration of Observational HIV Epidemiological Research Europe (COHERE) study group. Response to combination antiretroviral therapy: variation by age. *AIDS*. 2008;22(12):1463-1473.
66. Nogueras M, Navarro G, Anton E, et al. Epidemiological and clinical features, response to HAART, and survival in HIV-infected patients diagnosed at the age of 50 or more. *BMC Infect Dis*. 2006;6:159.
67. Bosch RJ, Bennett K, Collier AC, et al. Pretreatment factors associated with 3-year (144-week) virologic and immunologic responses to potent antiretroviral therapy. *J Acquir Immune Defic Syndr*. 2007;44(3):268-277.
68. Fahey JL, Taylor JM, Detels R, et al. The prognostic value of cellular and serologic markers in infection with human immunodeficiency virus type 1. *N Engl J Med*. 1990;322(3):166-172.
69. Giorgi JV, Lyles RH, Matud JL, et al. Predictive value of immunologic and virologic markers after long or short duration of HIV-1 infection. *J Acquir Immune Defic Syndr*. 2002;29(4):346-355.
70. Deeks SG, Kitchen CM, Liu L, et al. Immune activation set point during early HIV infection predicts subsequent CD4+ T-cell changes independent of viral load. *Blood*. 2004;104(4):942-947.
71. Giorgi JV, Hultin LE, McKeating JA, et al. Shorter survival in advanced human immunodeficiency virus type 1 infection is more closely associated with T lymphocyte activation than with plasma virus burden or virus chemokine coreceptor usage. *J Infect Dis*. 1999;179(4):859-870.
72. Hazenberg MD, Otto SA, van Benthem BH, et al. Persistent immune activation in HIV-1 infection is associated with progression to AIDS. *AIDS*. 2003;17(13):1881-1888.
73. Gandhi RT, Spritzler J, Chan E, et al. Effect of baseline- and treatment-related factors on immunologic recovery after initiation of antiretroviral therapy in HIV-1-positive subjects: results from ACTG 384. *J Acquir Immune Defic Syndr*. 2006;42(4):426-434.
74. Hunt PW, Martin JN, Sinclair E, et al. T cell activation is associated with lower CD4+ T cell gains in human immunodeficiency virus-infected patients with sustained viral suppression during antiretroviral therapy. *J Infect Dis*. 2003;187(10):1534-1543.
75. Neuhaus J, Jacobs DR, Jr., Baker JV, et al. Markers of inflammation, coagulation, and renal function are elevated in adults with HIV infection. *J Infect Dis*. 2010;201(12):1788-1795.
76. Valdez H, Connick E, Smith KY, et al. Limited immune restoration after 3 years' suppression of HIV-1 replication in patients with moderately advanced disease. *AIDS*. 2002;16(14):1859-1866.

77. Robbins GK, Spritzler JG, Chan ES, et al. Incomplete reconstitution of T cell subsets on combination antiretroviral therapy in the AIDS Clinical Trials Group protocol 384. *Clin Infect Dis*. 2009;48(3):350-361.
78. Lange CG, Lederman MM, Medvik K, et al. Nadir CD4+ T-cell count and numbers of CD28+ CD4+ T-cells predict functional responses to immunizations in chronic HIV-1 infection. *AIDS*. 2003;17(14):2015-2023.
79. Palella FJ, Jr., Gange SJ, Benning L, et al. Inflammatory biomarkers and abacavir use in the Women's Interagency HIV Study and the Multicenter AIDS Cohort Study. *AIDS*. 2010;24(11):1657-1665.
80. Rodger AJ, Fox Z, Lundgren JD, et al. Activation and coagulation biomarkers are independent predictors of the development of opportunistic disease in patients with HIV infection. *J Infect Dis*. 2009;200(6):973-983.
81. Centers for Disease Control and Prevention (CDC). Achievements in public health. Reduction in perinatal transmission of HIV infection--United States, 1985-2005. *MMWR Morb Mortal Wkly Rep*. 2006;55(21):592-597.
82. Panel on Treatment of HIV-Infected Pregnant Women and Prevention of Perinatal Transmission. Recommendations for use of antiretroviral drugs in pregnant HIV-1-infected women for maternal health and interventions to reduce perinatal HIV transmission in the United States. May 24, 2010:1-117. <http://aidsinfo.nih.gov/contentfiles/PerinatalGL.pdf>.
83. Vernazza PL, Troiani L, Flepp MJ, et al. Potent antiretroviral treatment of HIV-infection results in suppression of the seminal shedding of HIV. The Swiss HIV Cohort Study. *AIDS*. 2000;14(2):117-121.
84. Coombs RW, Reichelderfer PS, Landay AL. Recent observations on HIV type-1 infection in the genital tract of men and women. *AIDS*. 2003;17(4):455-480.
85. Quinn TC, Wawer MJ, Sewankambo N, et al. Viral load and heterosexual transmission of human immunodeficiency virus type 1. Rakai Project Study Group. *N Engl J Med*. 2000;342(13):921-929.
86. Tovnanabutra S, Robison V, Wongtrakul J, et al. Male viral load and heterosexual transmission of HIV-1 subtype E in northern Thailand. *J Acquir Immune Defic Syndr*. 2002;29(3):275-283.
87. Kayitenkore K, Bekan B, Rufagari J, et al. The impact of ART on HIV transmission among HIV serodiscordant couples. Paper presented at: XVI International AIDS Conference; August 13-18, 2006; Toronto, Canada. Abstract MOKC101.
88. Reynolds S, Makumbi F, Kagaayi J, et al. ART reduced the rate of sexual transmission of HIV among HIV-discordant couples in rural Rakai, Uganda. Paper presented at: 16th Conference on Retroviruses and Opportunistic Infections; February 8-11, 2009; Montreal, Canada. Abstract 52a.
89. Sullivan P, Kayitenkore K, Chomba E, et al. Reduction of HIV transmission risk and high risk sex while prescribed ART: Results from discordant couples in Rwanda and Zambia. Paper presented at: 16th Conference on Retroviruses and Opportunistic Infections; February 8-11, 2009; Montreal, Canada. Abstract 52bLB.
90. Granich RM, Gilks CF, Dye C, et al. Universal voluntary HIV testing with immediate antiretroviral therapy as a strategy for elimination of HIV transmission: a mathematical model. *Lancet*. 2009;373(9657):48-57.
91. Bunnell R, Ekwaru JP, Solberg P, et al. Changes in sexual behavior and risk of HIV transmission after antiretroviral therapy and prevention interventions in rural Uganda. *AIDS*. 2006;20(1):85-92.
92. Castilla J, Del Romero J, Hernando V, et al. Effectiveness of highly active antiretroviral therapy in reducing heterosexual transmission of HIV. *J Acquir Immune Defic Syndr*. 2005;40(1):96-101.
93. Wilson DP, Law MG, Grulich AE, et al. Relation between HIV viral load and infectiousness: a model-based analysis. *Lancet*. 2008;372(9635):314-320.
94. Worm SW, Sabin C, Weber R, et al. Risk of myocardial infarction in patients with HIV infection exposed to specific individual antiretroviral drugs from the 3 major drug classes: the data collection on adverse events of anti-HIV drugs (D:A:D) study. *J Infect Dis*. 2010;201(3):318-330.
95. Uy J, Armon C, Buchacz K, et al. Initiation of HAART at higher CD4 cell counts is associated with a lower frequency of antiretroviral drug resistance mutations at virologic failure. *J Acquir Immune Defic Syndr*. 2009;51(4):450-453.
96. Freedberg KA, Losina E, Weinstein MC, et al. The cost effectiveness of combination antiretroviral therapy for HIV disease. *N Engl J Med*. 2001;344(11):824-831.
97. Schackman BR, Goldie SJ, Weinstein MC, et al. Cost-effectiveness of earlier initiation of antiretroviral therapy for uninsured HIV-infected adults. *Am J Public Health*. 2001;91(9):1456-1463.
98. Mauskopf J, Kitahata M, Kauf T, et al. HIV antiretroviral treatment: early versus later. *J Acquir Immune Defic Syndr*. 2005;39(5):562-569.
99. Chen RY, Accortt NA, Westfall AO, et al. Distribution of health care expenditures for HIV-infected patients. *Clin Infect Dis*. 2006;42(7):1003-1010.
100. Willig JH, Abrams S, Westfall AO, et al. Increased regimen durability in the era of once-daily fixed-dose combination antiretroviral therapy. *AIDS*. 2008;22(15):1951-1960.
101. Bicanic T, Meintjes G, Rebe K, et al. Immune reconstitution inflammatory syndrome in HIV-associated cryptococcal meningitis: a prospective study. *J Acquir Immune Defic Syndr*. 2009;51(2):130-134.
102. Phillips P, Bonner S, Gataric N, et al. Nontuberculous mycobacterial immune reconstitution syndrome in HIV-infected patients: spectrum of disease and long-term follow-up. *Clin Infect Dis*. 2005;41(10):1483-1497.
103. Velasco M, Castilla V, Sanz J, et al. Effect of simultaneous use of highly active antiretroviral therapy on survival of HIV patients with tuberculosis. *J Acquir Immune Defic Syndr*. 2009;50(2):148-152.
104. Abdool Karim SS, Naidoo K, Grobler A, et al. Timing of initiation of antiretroviral drugs during tuberculosis therapy. *N Engl J Med*. 2010;362(8):697-706.
105. Kaplan JE, Benson C, Holmes KH, et al. Guidelines for prevention and treatment of opportunistic infections in HIV-infected adults and adolescents: recommendations from CDC, the National Institutes of Health, and the HIV Medicine Association of the Infectious Diseases Society of America. *MMWR Recomm Rep*. 2009;58(RR-4):1-207.
106. Lu M, Safren SA, Skolnik PR, et al. Optimal recall period and response task for self-reported HIV medication adherence. *AIDS Behav*. 2008;12(1):86-94.

107. Simoni JM, Kurth AE, Pearson CR, et al. Self-report measures of antiretroviral therapy adherence: A review with recommendations for HIV research and clinical management. *AIDS Behav.* 2006;10(3):227-245.
108. Bisson GP, Gross R, Bellamy S, et al. Pharmacy refill adherence compared with CD4 count changes for monitoring HIV-infected adults on antiretroviral therapy. *PLoS Med.* 2008;5(5):e109.
109. Kalichman SC, Amaral CM, Cherry C, et al. Monitoring medication adherence by unannounced pill counts conducted by telephone: reliability and criterion-related validity. *HIV Clin Trials.* 2008;9(5):298-308.
110. Moss AR, Hahn JA, Perry S, et al. Adherence to highly active antiretroviral therapy in the homeless population in San Francisco: a prospective study. *Clin Infect Dis.* 2004;39(8):1190-1198.
111. Hunt PW, Brenchley J, Sinclair E, et al. Relationship between T cell activation and CD4+ T cell count in HIV-seropositive individuals with undetectable plasma HIV RNA levels in the absence of therapy. *J Infect Dis.* 2008;197(1):126-133.
112. Choudhary SK, Vrisekoop N, Jansen CA, et al. Low immune activation despite high levels of pathogenic human immunodeficiency virus type 1 results in long-term asymptomatic disease. *J Virol.* 2007;81(16):8838-8842.
113. Egger M. Outcomes of ART in resource-limited and industrialized countries. Paper presented at: 14th Conference on Retroviruses and Opportunistic Infections; 2007; Los Angeles, CA. Abstract 62.
114. Wolbers M, Bucher HC, Furrer H, et al. Delayed diagnosis of HIV infection and late initiation of antiretroviral therapy in the Swiss HIV Cohort Study. *HIV Med.* 2008;9(6):397-405.
115. Grigoryan A, Hall HI, Durant T, et al. Late HIV diagnosis and determinants of progression to AIDS or death after HIV diagnosis among injection drug users, 33 US States, 1996-2004. *PLoS One.* 2009;4(2):e4445.
116. Centers for Disease Control and Prevention (CDC). Late HIV testing - 34 states, 1996-2005. *MMWR Morb Mortal Wkly Rep.* 2009;58(24):661-665.
117. Branson BM, Handsfield HH, Lampe MA, et al. Revised recommendations for HIV testing of adults, adolescents, and pregnant women in health-care settings. *MMWR Recomm Rep.* 2006;55(RR-14):1-17.